

Living with COVID: management of COVID-19 for New Brunswick Long Term Care Homes

Department of Social Development

November 2022

Working Group Members/ Acknowledgements

Natalie Callan (Document Writer)

Nurse Consultant
Department of Social Development

Brittany Jones (Document Writer)

Nurse Consultant
Department of Social Development

Serena Ann Bradford

Manager
Department of Social Development

Mathieu Chalifoux

Chief Epidemiologist
Department of Health

Chris Cohoon

Infection, Prevention and Control Consultant
Department of Social Development

Dr. Yves Leger

Acting Chief Medical Officer of Health
Public Health New Brunswick

Tasha Rossignol

Nurse Consultant
Department of Social Development

Liena Roussel

Manager
Department of Social Development

Dr. Andrew Salmon

Medical Officer of Health
Public Health New Brunswick

Alex Taylor

Manager
Department of Social Development

Joyce Walker-Hayley

Senior Program Advisor
Department of Health

Mary Williams

Acting Director, Home Care Unit
Department of Health

Purpose

This document is a revised version of the Living with COVID: management of COVID-19 for New Brunswick Long Term Care homes, which was published in July 2022 and replaced the following two documents:

- Living with COVID: General LTC Guidance (for homes NOT in outbreak)
- Living with COVID: Case & Outbreak Management in LTC Homes

In addition to the information that was contained in the documents listed above, this document includes additional resources such as key contact information and information on COVID-19 prevention measures. The document has also been re-organized in the following sections for ease of use:

1. [Prevention](#)
2. [Case Management](#)
3. [Outbreak Management](#)
4. [Appendices](#)

This is a *living document* (continually edited and updated) and will be revised as new information becomes available. **Unless otherwise indicated, information contained in this guide applies to both Adult Residential Facilities (ARFs) and Nursing Homes (NHs) which will further be referred to as long-term care (LTC) Homes. Any site-specific information will be identified as Nursing Home (NH) or Adult Residential Facility (ARF) in the heading.**

Adult Residential Facilities with a license for 10 or fewer beds: An ARF with 10 or fewer residents often is structured and functions similar to a household. Moving forward, they will self-manage individual positive COVID-19 cases within their facility, or situations involving 2 or more positive individuals. Outbreaks will no longer be declared within these facilities. Best practice recommendations for case and situation management in this setting still apply and are listed in [Appendix 4.1](#).

Resources and Contact Information

Websites				
COVID-19 Test Registration Form		COVID-19 Testing (gnb.ca)		
General GNB COVID-19 Resources		Guidance and Support (gnb.ca)		
Vaccination Survey for Employees and Residents		https://survey.alchemer-ca.com/s3/50129571/COVID19VACCINATION		
Emails and Phone Numbers				
SD PPE Team		PPETeam/EquipeEPI@GNB.CA		
SD COVID Response Team (SD CRT)		SD_CRT@gnb.ca		
COVID-19 Vaccine Orders (for NHs)		DHvacclg@gnb.ca		
Extra Mural Program- EMP Care Coordination Centre		1-844-982-7367		
Public Health Numbers		(see table below)		
Public Health				
Regions1-7 Public- Health Notifiable Disease Reportable Events Posters.pdf (gnb.ca)				
Region		During Business Hours (8:15'am-4:30pm Mon-Fri)	After Business Hours (EMERGENCY only)	Fax Numbers
Central	Zone 3 (Fredericton)	CDFredVal@gnb.ca (506) 444-5905	(506) 453-8128	(506) 444-4877
South	Zone 2 (Saint John)	ComDisjPH@gnb.ca (506) 658-5188	(506) 658-2764	(506) 643-7894
East	Zone 1 (Moncton)	(506) 856-3220	(506) 856-2004	(506) 856-3544
	Zone 7 (Miramichi)	(506) 778-6104 / 778-6102		(506) 778-6756
North	Zone 4 (Edmundston)	(506) 735-2626	(506) 789-2428	(506) 735-2340
	Zone 5 (Campbellton)	(506) 790-4769		(506) 789-2349
	Zone 6 (Bathurst)	(506) 547-2062		(506) 394-3858

¹ Zone 2 business hours start at 08:30

COVID-19 MANAGEMENT FOR NB LTC HOMES DOCUMENT OVERVIEW

SECTION	KEY INFORMATION	SUPPORTING INFO
<p><u>1. PREVENTION</u> Practices and measures to be in place to prevent introduction and transmission of COVID-19.</p>	<ul style="list-style-type: none"> • Immunization 1.1 • PPE 1.2 • Screening 1.3 • Testing 1.5 	<ul style="list-style-type: none"> • Hand Hygiene 4.3 • Cleaning & disinfecting 4.5 • Re-testing 4.6 • Visitation 4.7
<p><u>2. CASE MANAGEMENT</u> Measures for individual COVID-19 cases.</p>	<ul style="list-style-type: none"> • Work Exclusion 2.1 / Isolation 2.2 • Enhanced Work Precautions 2.1.1 • Treatment 2.3 	<ul style="list-style-type: none"> • COVID-19 positive household member 2.1.2 • Admission 4.9 •
<p><u>3. OUTBREAK MANAGEMENT</u> Measures to manage a declared COVID-19 outbreak.</p>	<ul style="list-style-type: none"> • General Outbreak Principles 3.1 • Outbreaks that DO NOT include at least 1 resident 3.2 • Outbreaks that included at least 1 resident 3.3 	<ul style="list-style-type: none"> • Visitation during an outbreak 4.8 • Admission 4.9
<p><u>4. APPENDICES</u> Additional resources that may be useful.</p>	<ul style="list-style-type: none"> • Management of one or more cases of COVID-19 in ARFs with 10 or fewer residents 4.1 • Posters 4.2 • Hand Hygiene 4.3 • PPE Table 4.4 • Cleaning & disinfecting 4.5 	<ul style="list-style-type: none"> • Re-testing 4.6 • Visitation 4.7 • Visitation during an outbreak 4.8 • Admission 4.9 • POCT throat - nares swabbing guidance 4.10

1. Prevention

1.1. COVID-19 Immunization

1.1.1. Staff Immunization

Definition of up to date COVID-19 Vaccination (for vulnerable sector): primary series plus a booster in the last 6 months.

Employees/volunteers, within the vulnerable sector, who are not up to date for their COVID-19 vaccination can work on the condition they follow all [mitigation measures](#) required by their respective employer, as determined by the NB Infection Prevention and Control Committee. If/when an employee decides to get vaccinated, mitigation measure may be removed once they are considered up to date.

Evaluation of mitigation measures will be ongoing. To promptly implement any change in mitigation measures based on epidemiology risk, employers must maintain a list² of the COVID-19 vaccination status for all employees and all volunteers. Homes will continue to demonstrate compliance through the routine process of inspection.

[Mitigation Measures for staff who are not up to date with their COVID-19 Vaccination \(including unvaccinated\)](#)

Homes are required to develop and implement policy aligning with the mitigation measures listed below.

- 1. Masking:** mask must be worn at all times, unless eating or drinking.
- 2. Screening:** Passive symptom screening is required at the beginning of every shift.

Important Note: masking and screening are currently required for all employees (even those considered to have an up to date COVID-19 immunization status). However, should policies change in the future (e.g. masking policy), the mitigation measures listed above will be considered a minimum requirement for employees whose vaccination against COVID-19 is not considered up to date.

1.1.2. Resident Immunization

New Brunswick recommends that everyone should receive a complete COVID-19 vaccination series and a booster (first or second) as soon as they become eligible to protect themselves from COVID, including residents of long-term care facilities. Operators/staff of ARFs and NHs should coordinate immunization for residents as soon as they become eligible for boosters. **Residents who are determined to be eligible for admission to an ARF or NH are to be admitted, regardless of their COVID-19 vaccination status.**

² The home is not to retain a copy of the actual personal vaccination documentation of employees and volunteers.

1.2. Personal Protective Equipment (PPE)

It is recommended that LTC homes have a minimum amount of PPE on hand, please refer the table below for the recommended amount for a 72-hour period where outbreak measures would be in place.

Emergency PPE Recommended for 72 Hour Period

Number of Beds	Quantity Recommended for 72 Hours			
	Face Masks	Isolation Gowns	Face Shields	Gloves
10	300	300	300	1,050
30	900	900	900	3,150
60	1,800	1,800	1,800	6,300
100	3,000	3,000	3,000	10,500

1.2.1. Staff PPE

It is imperative that PPE be used correctly to protect staff and residents. Improper use of PPE can lead to rapid transmission throughout facilities. Refer to [PPE Table](#) for an overview of PPE that is used for the prevention of spread of COVID-19.

Please note additional use of PPE may be directed by PH in certain situations or based on a point of care risk assessment.

- a) **Masks:** continuous mandatory masking using medical grade masks in any area where staff may come in contact with a resident. For staff who are not considered up to date for their COVID-19 vaccination, they must follow the masking guidance as per the [risk mitigation measures](#) and wear a mask at all times unless eating or drinking. Additional use of PPE should be based on PCRA (point of care risk assessment) and/or as directed by MOH in certain situations.
- b) **Eye protection:** based on PCRA, optional for continuous use, unless required during an outbreak as per outbreak management guidelines.

1.2.2. Resident Masking

Masks should be made available to residents if they wish to wear one.

1.3. Staff Screening

A facility specific policy for workplace health that must include direction to not work while ill (form of passive screening). Staff should stay home when ill and get tested. Using an established protocol when staff call in ill or become ill during their shift will ensure a healthy and productive workforce. Screening questions should include: is there anyone sick at home with COVID symptoms?" If yes, staff should follow the measures listed in section [2.1.2 Staff with Household Member who has tested positive for COVID-19 or has recent ILI symptoms](#)). Household members of staff working in vulnerable settings should monitor for symptoms and get tested if symptomatic to provide additional protective measures to residents.

1.4. Resident Assessments

Residents should be assessed as clinically indicated and as per facility specific policy.

Guidance for ARF with 10 or fewer residents:

- Monitor resident's symptoms and ability to cope with illness daily. i.e., Fever, cough or difficulty breathing, diarrhea, fatigue; and stability, alertness, intake.
- Refer to [Caring for Someone with Covid-19](#).
- If there are concerns related to the health of the resident, contact their primary care provider. If resident does not have a primary care provider, or they are unavailable, contact EMP, **EMP Care Coordination Center 1-844-982-7367**.
- Watch for signs that indicate immediate help is needed. Call 911* if you notice:
 - significant difficulty breathing;
 - chest pain or pressure;
 - new onset of confusion; or
 - difficulty waking up.

*Make sure you let 911 and the hospital know that you or the person you're caring for has symptoms or has tested positive for COVID-19.

1.5. Testing

1.5.1. Staff Testing

PCR testing is recommended for staff who have a positive POCT, meet the [symptom checker criteria](#), or as directed by MOH as necessary. Please see [Appendix: Retesting](#) for staff who previously tested positive (PCR/POCT).

Staff can book an appointment for Polymerase Chain Reaction (PCR testing) at [COVID-19 Testing](#) . Alternatively, if a staff member is unable to book an appointment online, they may also do so by calling 811 and **identify themselves as staff of LTC**.

Employees/volunteers must immediately notify their employer if they receive a positive PCR test result on MyHealthNB or have received a positive POCT and are awaiting PCR confirmation. **Recommendation:** all employees should create a MyHealthNB account. This can be done by registering online: <https://id.qnb.ca/registration>

1.5.2. Resident Testing

Only [symptomatic](#) residents should be referred for PCR testing, unless directed by MOH. Please see [Appendix: Retesting](#) for residents who previously tested positive (PCR/POCT). If resident tests positive and outbreak is declared, follow guidance in section [3.3.2: Resident Testing during an outbreak that includes at least one resident](#) .

- **For ARFs:** contact the EMP care coordination center (1-844-982-7367) to arrange PCR testing of symptomatic residents. If mass testing is required, it will be ordered by PH
- **For NHs:** complete the combined referral and requisition form, collect sample for PCR testing and send to laboratory for processing.
- **NOTE:** a positive result from a rapid POCT test performed by trained staff or observed by a health care professional may be used to hasten access to Paxlovid.

1.6. Public Health Reporting

- 1.6.1. ARFs and NHs with greater than 10 residents:** the first employee/staff person or resident testing positive with POCT/PCR shall be reported to Public Health within 1 hour. All deaths of COVID positive residents or staff who work or live in an LTC facility are required to be reported to Public Health. The LTC home must inform Public Health whether COVID was a primary or contributing factor to cause of death as per the attending physician. If in a declared outbreak, report subsequent cases once daily³ by line list via email or fax to PH CD Team to fulfill reporting requirements, as per the Public Health Act.
- 1.6.2. ARFs with 10 or fewer residents:** LTC home must report COVID hospitalizations and deaths to PH. Reporting should be done during [PH regular business hours](#).

1.7. Reporting to Social Development

- 1.7.1. SD COVID Response Team:** if Public Health declares an outbreak, please send an email to the centralized team at SD_CRT@gnb.ca as soon as possible with the following details: Name of home, date your outbreak was declared, and any questions you may have.
- 1.7.2. Reporting incidents as per ARF and NH standards.** In addition to reporting to PH and to the SD COVID Response Team, ARFs and NHs are to continue reporting incidents as per the process and requirements from their respective standards.

³ RMOH will specify time of day that line lists are to be received.

2. Case Management

2.1. Staff Work Exclusion

<p>If tested & positive (POCT / PCR)</p>	<ul style="list-style-type: none"> Exclude from work for 5 days (test date is day 0) and until symptoms are improving and fever free x 24hrs. Required to use Enhanced Work Precautions for an additional 5 days after returning to work.
<p>If Critical Staffing Shortage</p>	<ul style="list-style-type: none"> Exclude from work if symptomatic and until symptoms are improving and fever free x 24hrs. Can return to work using Enhanced Work Precautions for the balance of 10 days from symptom onset or positive test result if no symptoms.
<p>If staff refuses to get tested & symptoms meet symptom checker criteria</p>	<ul style="list-style-type: none"> Exclude for 5 days and until symptoms are improving (and must be without fever for 24 hrs). Can return to work with Enhanced Work Precautions for the balance of 10 days from symptom onset or positive result (if asymptomatic).

2.1.1. Enhanced Work Precautions

- Wear a well fitted medical grade mask.
- Perform [hand hygiene](#) frequently.
- Maintain physical distancing and limit non-essential contact with other staff, and residents (as able).
- Avoid meeting spaces and lunchrooms.
- Eating/drinking must be performed in a private area. If you cannot find a place to eat or drink alone, ensure there is two metre distancing from others while your mask is off or consider staggering lunch hour for staff who are positive and negative etc.
- Use washrooms within the organization which are the most frequently cleaned if a designated washroom is not possible.
- Employees should also clean high touch areas after use.

2.1.2. Staff with Household Member who has tested positive for COVID-19 or has a recent ILI symptoms

- If staff is symptom free, no work exclusion is required
- [Enhanced Work Precautions](#) for staff member X 10 days (see [note](#) below)
- Day 5 PCR [test](#) (test should occur regardless of the employee's ability to isolate from the household case)

NOTE: enhanced work precautions are in place for staff who have household contacts as well as staff returning to the workplace while positive for COVID-19. The risk of transmission varies in both situations, the higher risk situation being the staff who is a household contact. Because of the difference in risk, it is important that breaks be staggered for staff on enhanced work precautions and all enhanced work precautions followed.

2.2. Resident Isolation

Residents with any [COVID-19 symptoms](#) should be placed on **contact droplet precautions**.

2.2.1. COVID-19 test result - NEGATIVE:

If the test result is negative it is important to ensure appropriate clinical assessment to determine if the resident may have an alternate condition or infection other than COVID-19 which needs further medical intervention or assessment.

2.2.2. COVID-19 test result - POSITIVE:

- Isolate for 5 days after testing.
- Remove from isolation after day 5 with improving symptoms and minimum 24 hrs without fever.
- For an additional 5 days, [extra measures](#) are in place for resident.

Extra Measures:

- resident to wear a mask (as able)
- physically distance
- eat meals distanced from other residents.

2.3. Treatment - Paxlovid™

Primary care providers can prescribe nirmatrelvir/ritonavir (PAXLOVID™) for COVID-19 positive residents who are [eligible](#). Eligibility form can be found here:

<https://www2.gnb.ca/content/dam/gnb/Departments/h-s/pdf/en/NBDrugPlan/PaxlovidEligibilityForm.pdf>

Please refer to [Section 2.14](#) of this document for further detail on Paxlovid.

3. Outbreak Management

This section of the document is intended for use by operators and managers of LTC homes to guide their decision-making related to outbreak response.

Public Health remains committed to supporting LTC homes in managing COVID-19.

- Public Health will lead and manage outbreaks⁴ within LTC homes if the [outbreak includes at least 1 resident](#).
- For LTC homes who have [outbreaks that do not include any residents](#), the home will self-manage unless directed by Public Health.

3.1. General Outbreak Management Principles⁵

- a) Public Health will declare the start and end to outbreaks.
- b) Resident assessment: daily active monitoring of residents.
- c) Staff screening: passively screen before every shift and self monitor throughout shift.
- d) Work exclusion for staff who meet [symptom checker criteria](#).
- e) Enhance cleaning and disinfection of staff areas (break/lunchrooms, workstations etc)
- f) Stringent adherence to masking and eye protection at all times for staff working on affected unit/area.
- g) Ensure staff break/lunch areas do not promote transmission – limit capacity, ensure adequate distancing and barriers, if possible, between staff, wipes for disinfection etc.
- h) **Reporting Guidance:** if an outbreak is confirmed, Public health will communicate this to the home. Subsequent cases should be reported in a daily line list that will be sent to Regional Public Health at an agreed upon time in the day.

It is important for homes to note that not all outbreaks are the same. Therefore, guidance will vary depending on each situation and circumstances. Direction may be adjusted by Public Health / the MOH as needed based on the result of their risk assessment and applicable factors. If homes are unsure why certain directives might be given to them, they should seek to have their questions clarified.

3.2. Outbreak that does NOT include a resident

3.2.1. Role of PH

Supportive (including declaration of beginning and end of outbreak); facility self manages using [general outbreak management principles](#).

3.2.2. End of Outbreak

End 10 days after last exposure.

3.3. Outbreak that includes at least one resident

3.3.1. Role of PH (outbreak that includes at least one resident)

Active including:

- Initial touchpoint with facility to review the situation, declare the outbreak and adjust/clarify the above guidance as well as any additional guidance required.

⁴ COVID-19 outbreaks are defined as 2 laboratory confirmed cases in either staff or resident where transmission on site has not been ruled out. Outbreaks are declared by MOH.

⁵ Outbreak management principles apply to residents and staff of affected unit, unless otherwise directed by MOH.

- Receive daily line lists to maintain situational awareness.
- Answer questions from facility.
- Involved in determining risk level of exposures.
- Declare outbreak over.

3.3.2. Resident Testing during an outbreak that includes at least one resident

- Once an outbreak has been declared, testing of all symptomatic residents may not be required.
- Cases (probable) may be identified and managed based on presence (one or more) new or worsening symptoms.
- One round of mass testing may be prescribed by the MOH. The MOH will determine the timing of such testing if required.
- Testing may be prescribed for the purpose of determining eligibility to treatment (i.e.: Paxlovid)
- Refer to [Appendix: Re-testing](#) for residents who have recently tested positive for COVID-19.
- Additional testing may be required to rule out co-infections with other respiratory illnesses.

3.3.3. Control measures to be implemented during an outbreak that includes at least one resident:

- Restrict residents to affected wings/units
- Group activities can continue provided small, consistent groups are maintained. Masking and physical distancing for residents as able.
- General visitation restricted except for designated support persons or when residents are palliative, see [Appendix: Visitation during an outbreak](#) for details on visitation during an outbreak. [General visitation](#) may occur *during an outbreak* if facility has implemented control measures and is able to adequately function in outbreak management as determined by PH.

3.3.4. End of Outbreak

10 days after last high-risk exposure (as determined by PH).

4. Appendices

4.1. Management of one or more cases of COVID-19 in ARFs with 10 or fewer residents

Best practice recommendations for case and situation management in this setting still apply and are listed below. However, it is recognized that ability to implement may be limited by staffing levels, facility size, or capacity of residents:

Testing	<ul style="list-style-type: none"> • Symptomatic residents and staff only, unless prescribed by guidance or to access treatment. • Once one case of COVID-19 is identified, no need to test others with symptoms unless eligible for treatment. • If negative for Covid-19, additional testing may be needed for other pathogens (i.e. Influenza, RSV), or to access prophylactic Tamiflu for contacts.
Case and Symptomatic Resident Isolation	<ul style="list-style-type: none"> • Cases and symptomatic residents isolate separately, and apart from other residents, as best as possible for 5 days and until fever free for 24 hrs if facility space allows. • A symptomatic person or positive case can eat in their room; or consider having different dining shifts to increase separation between residents at table.
Daily Care	<ul style="list-style-type: none"> • Monitor resident's symptoms and ability to cope with illness daily. i.e., Fever, cough or difficulty breathing, diarrhea, fatigue; and stability, alertness, intake. <ul style="list-style-type: none"> ○ Refer to Caring for Someone with Covid-19. • If there are concerns related to the health of the resident, contact their primary care provider. If resident does not have a primary care provider, or they are unavailable, contact EMP, EMP Care Coordination Center 1-844-982-7367. • Watch for signs that indicate immediate help is needed. Call 911* if you notice: <ul style="list-style-type: none"> ○ significant difficulty breathing; ○ chest pain or pressure; ○ new onset of confusion; or ○ difficulty waking up. • *Make sure you let 911 and the hospital know that the resident has symptoms or has tested positive for COVID-19.
Washroom	<ul style="list-style-type: none"> • Designate separate washroom for case if possible. • If shared with others, try to clean between users or as often as possible each day.
Cleaning	<ul style="list-style-type: none"> • Per section 4.5, as able.
Staff PPE	<ul style="list-style-type: none"> • Resources permitting, and where possible, contact/droplet precautions are recommended with cases for their 5 days of isolation as well as with any other symptomatic residents. • Mask and eye protection are also recommended for 5 days after isolation.
Group activities	<ul style="list-style-type: none"> • Active daily monitoring. Only asymptomatic participate. • On site, with masking and distancing in place as much as possible. Provide hand sanitizer and encourage cough/sneeze etiquette.
Visitors	<ul style="list-style-type: none"> • DSP only, or palliative visitation permitted.
End of additional precautions	<ul style="list-style-type: none"> • 5 days after last case identified

4.2. Posters/Printable resources

If you require posters for donning and doffing PPE or PCRA (point of care risk assessment), please email SD_CRT@gnb.ca.

For NHs - these resources are available on iTacit.

4.3. Hand Hygiene

Hand hygiene is the single best way to prevent spread of infection. It is estimated that 80% of common infections such as the cold and flu are spread by contaminated hands. Use of an alcohol-based hand rub (ABHR) with between 70-90% ethyl alcohol is best practice for ARFs and NHs. Facilities should continue with a comprehensive hand hygiene program that includes all departments with involvement from staff, residents, and families.

Hands must be cleaned at the point of care and it is crucial that hand hygiene is performed at these **4 critical moments:**

1. Before initial resident/resident environment contact.
2. Before aseptic procedure.
3. After body fluid exposure risk.
4. After resident/resident environment contact.

Personal hand hygiene should also be performed:

- Upon entering facility
- Before assisting residents with meals
- Before and after preparing food
- Before and after eating meals
- Before and after smoking/vaping
- After using the rest room
- After blowing your nose, coughing or sneezing

Please note: if there is visible soiling or exposure to bodily fluids, hands should be washed with soap and water. Soap and water should also be used after using the toilet.

4.4. PPE Table

ALL PPE MUST BE CHANGED IF IT BECOMES SOILED, WET OR DAMAGED		
Type	Description	Tips
Masks <i>**continuous use**</i>	Face masks (ASTM level 1) provide a physical barrier that help prevent the transmission of the virus from person to person by blocking large particle respiratory droplets propelled by coughing or sneezing	Must be changed upon exit of an isolation room, UNLESS providing sequential care between COVID POSITIVE residents <u>OR</u> providing sequential care between NON COVID residents.
N95 respirators	Respiratory protective device designed to have a very close facial fit and efficient filtration of airborne particles.	Edges of the respirator are designed to form a seal around the nose and mouth.
Eye Protection	Eye protection (e.g. goggles, face shields etc.) protect the mucous membranes of the eyes during case/probable case/suspect case care or activities likely to generate splashes or sprays of body fluids including respiratory secretions. Prescription eyeglasses alone are not adequate protection against respiratory droplets.	Must be changed (or cleaned if using non-disposable eye protection) upon exit of an isolation room, UNLESS providing sequential care between COVID POSITIVE residents <u>OR</u> providing sequential care between NON COVID residents.
Gloves	Disposable single use gloves should be worn when in direct contact with the ill person, cleaning contaminated surfaces, and handling items soiled with body fluids, including dishes, cutlery, clothing, laundry, and waste for disposal.	<ul style="list-style-type: none"> • Must be changed between each resident. • Not a substitute for hand hygiene; staff must perform hand hygiene before putting on and taking off gloves. • Not for continuous use outside of an isolation room. • Gloves should be removed, hand hygiene performed, and new gloves applied when they become soiled or torn during care. • Double gloving is not necessary.
Gown	Used to protect your clothing and skin from becoming contaminated/soiled from: splashes/sprays of blood, body fluids, secretions or excretions, contact with soiled/contaminated items/surfaces.	<ul style="list-style-type: none"> • Must be changed between each resident. • Not for continuous use outside of an isolation room. • Must be tied at the neck and the back. If clothing exposed underneath, two gowns should be worn, first covering the back (tied in front), second covering front (tied in back).
Donning & Doffing	<p>Proper donning and doffing (putting on and taking off) of PPE is key in preventing the spread of COVID-19 from one individual to another.</p> <p>When removing PPE (doffing):</p> <ul style="list-style-type: none"> • PPE below the neck: gown and gloves, are removed <i>in the room (close to the door)</i>. • PPE above the neck: eye protection and masks, are changed <i>outside of the room</i>. 	

4.5. Cleaning and Disinfection

Continue regular cleaning and disinfecting of all general surfaces. Increasing the frequency of cleaning and disinfecting high-touch surfaces is significant in controlling the spread of viruses, and other microorganisms. All surfaces, especially those general surfaces that are frequently touched, such as doorknobs, handrails, etc., should be cleaned and disinfected at least twice daily and when soiled. Use an environmentally acceptable [cleaning and disinfecting product for COVID-19](#).

Cleaning & Disinfection	
Area and Description	Frequency (min)
Surfaces: in resident rooms and central areas	
Horizontal (e.g. tables)	Twice Daily (and when soiled)
High touch (e.g. telephone, bedside table, overbed table, chair arms, call bell cords or buttons, door handles, light switches, bedrails, handwashing sink, bathroom sink, toilet and toilet handle and shower handles, faucets or shower chairs, grab bars, outside of paper towel dispenser.)	
Low touch (e.g. shelves, bedside chairs or benches, windowsills, headwall units, overbed light fixtures, message or white boards, outside of sharps containers).	Once Daily (and when soiled)
Resident Care Equipment	
E.g. BP cuffs, electronic thermometers, oximeters, stethoscope etc. Hospital grade disinfectant (e.g., cleaner and disinfectant wipes) using the recommended contact time should be used to clean and disinfect smaller resident care equipment. Important Note: Ensure all staff responsible for utilizing resident care equipment is adhering to required cleaning and disinfection practices.	After Each Use
Single Use Devices: discard in a waste receptacle after a single use on one resident. Single person/resident devices: discard after use with one resident (may be more than one use).	
Outside Resident Room (e.g. surfaces that are touched by or in contact with staff)	
Computer carts and/or screens, medication carts, charting desks or tables, computer screens, telephones, touch screens, chair arms. Staff should ensure that hands are cleaned before touching the above-mentioned equipment.	Once Daily (minimum) and when soiled.
Floors and Walls: Kept visibly clean and free of spills and debris.	
Terminal Cleaning and Disinfecting	
Items that cannot be appropriately cleaned and disinfected should be discarded upon resident transfer or discharge. Toilet brushes, unused toilet paper and other disposable supplies should be discarded. Privacy curtains should be removed and laundered upon a resident's discharge or transfer.	After discharge, transfer or discontinuation of contact droplet precautions.
Ensure all staff responsible for environmental cleaning adhere to required cleaning and disinfection practices.	

4.6. Retesting - recommendations for individuals who have previously tested positive for COVID-19 (PCR/POCT) in LTC

Recommendations are based on the length of time since the positive COVID-19 result:	
If more than 90 days	PCR testing is recommended if individual is symptomatic. If resident/staff are positive, follow guidance outlined in this document.
Between 30 and 90 days	<p>For individuals experiencing new or worsening symptoms, testing (POCT) is recommended. Rationale: POCT is recommended during this time to reduce the risk of false positives from a recent previous infection when using a PCR test.</p> <ul style="list-style-type: none"> • Staff/Residents should have two POCTs taken between 24-48 hours apart. • Note that if the first test is positive, the second test is not required. • Should symptoms worsen or new symptoms appear, the test should be repeated. <p>Individuals currently symptomatic, or who have tested positive, should follow the guidance on isolation and work exclusion included in this document.</p> <ul style="list-style-type: none"> • If the test(s) is/are negative: the individual may cease isolation/work exclusion, if asymptomatic. If they become symptomatic <u>or</u> their symptoms worsen, they must isolate/work exclude and get retested.
Less than 30 days	<p>No additional PCR testing is generally recommended. However, should a risk of exposure and reinfection be of concern, POCT use may be considered. No additional COVID-19 isolation or work exclusion requirements. For new or worsening symptoms, the individual may be feeling sick due to another reason.</p> <ul style="list-style-type: none"> • Residents should be clinically assessed to determine any other possible cause of symptoms and be placed on the appropriate precautions. • Staff should stay home while ill, as per site specific workplace health policy.

4.7. General Visitation

Each facility must have a visitation policy which includes:

- No visiting when ill, signage at door stating not to enter if unwell
- Visitors should be provided information outlining safe visiting practices, including frequent hand hygiene and respiratory etiquette with each visit, and postponing a visit when ill.
- Visitors must wear a well-fitting medical grade face mask while travelling through facility or in common areas. Mask may be removed in resident room if resident agrees. Eye protection is not required for visitors unless the resident they are visiting is on contact and droplet precautions.
- There is no age restriction related to visitation within an LTC facility.
- There is no limit on visitation capacity at one time.
- There is no requirement for visitors to provide proof of vaccination or medical exemption.

During an outbreak, certain visitation continues to be permitted, please see [Visitation During an Outbreak](#).

4.8. Visitation During an Outbreak

VISITATION GUIDANCE DURING OUTBREAK	
For units/facilities which are in outbreak certain types of visitation are permitted as outlined below.	
Designated Support Persons (DSPs)	Permitted into affected units/facilities . These individuals are to be informed that an outbreak is occurring and to visit one resident within the facility only and not go from room to room. DSPs must follow the same infection control measures as staff, including the appropriate use of PPE during visits as indicated. If resident is not on isolation precautions DSP must wear medical grade mask and eye protection continuously as required as part of outbreak management.
Palliative⁶ Visitation	<ul style="list-style-type: none"> • In collaboration with the home, a physician or nurse practitioner determines if the resident's condition is considered end-of-life based on current clinical assessment findings. • When death is anticipated as imminent, palliative visitation may occur, with a visitation plan developed in consultation with the care team of the home. • Resident/POA/substitute decision maker may have a list of 10 visitors • Homes should develop a safe visitation plan for any palliative visitation which will occur. Safe visiting interventions may include: <ul style="list-style-type: none"> ○ Visitors understand the risk of exposure to COVID-19 (for self and others); ○ Visitors follow all related home policies and public health measures in place within the home; and ○ Visitors remain vigilant in protecting themselves and others while on site, including: use of continuous medical grade mask and eye protection appropriate hand hygiene, following isolation precautions as required.

⁶ For the purposes of this document, palliative residents, are defined as individuals whose condition is considered end-of-life and death is anticipated as imminent.

4.9. Admissions & Readmissions (return from hospital)

Individuals should not be unreasonably denied admission or readmission. When a possible risk of COVID has been identified, no admissions, readmission or transfer will be considered if either of the following criteria are met:

- Inability to isolate the resident (either availability of private room and bathroom, cognitive impairments that would impede isolation or other operational considerations), applicable to situations where a resident requires to be isolated upon admission
- Critical Staffing at the receiving NH/ARF facility.

It is recognized some hospitalized patients, determined as medically stable / ready for discharge, have been unable to be newly admitted or returned to their Nursing Home / Adult Residential Facility (NH/ARF) due to COVID. The situations where this may occur include:

- The NH/ARF has a current COVID outbreak
- The patient has had a possible COVID exposure during their hospitalization (in a room, on a unit with a COVID outbreak), OR,
- Is COVID positive and medically stable/discharged; and must continue isolation period post-discharge.

Admissions and readmissions to an affected unit/wing during an outbreak in the ARF/NH are not recommended unless the person was determined to be COVID positive within the last 90 days. Should it be necessary to admit residents during the outbreak the following factors apply:

- The outbreak is under control, as determined by PH.
- The resident's attending physician is in agreement based on the current health status of the resident.
- Appropriate accommodation is available.
- Staffing is appropriate to meet the needs of the resident.
- The resident or substitute decision-maker has given an informed consent.

4.10. Throat – Nasal Specimen Collection for COVID-19


The use of throat and nasal swabs has been recognized as effective for rapid antigen point of care testing (POCT) in long term care (LTC) homes where staff have been trained in this process. Evidence shows that although both nasal and throat/nasal methods are acceptable, combined throat/nasal samples provide better results for detection of COVID-19 and the Omicron variant.

How to do a throat/nasal swab for COVID-19:

1. **Advise resident not to eat, drink, vape, smoke, brush teeth or chew gum 30 minutes prior to testing.**
2. **Prepare testing supplies:** Wash hands before starting. Prepare the testing area and lay out the supplies you will need on a clean surface. POCT supplies needed are:
 - Tube rack
 - Testing device
 - Tube and cap
 - Buffer fluid
 - Swab (keep in wrapper until ready to test)
 - Personal Protective Equipment for the staff member
3. **Prepare the test according to package instructions:** Tests vary by brand. Note that some swabs are scored to break easily. Hold the swab with the thumb and index finger on the score line to prevent breakage when swabbing.
4. **Obtain Throat Swab:** Swab the tonsil area in the back of the throat for five seconds on each side (avoiding the tongue, teeth and lips).
5. **Obtain Nasal Swab:** Swab the nostrils using the same swab. Gently insert the swab about 1-2 cm into the first nostril, or until resistance is felt. Rotate five times, remove and repeat on the other side.
6. **Process the sample:** Place the swab in the tube. Follow the test kit instructions.
7. **Positive results do not require a confirmatory PCR test.** Refer to LTC Guidance regarding for next steps once results are displayed and testing is complete



(a) Throat technique



(b) Nose technique

References:

- EM/ANB Throat/Nose Specimen Collection.pdf
- Provincial Public Health Laboratory Nova Scotia [instructions-alternate-collection-throat-and-nares-covid.pdf](#)
- [Throat and Nares \(Nasal\) swabbing for Rapid Antigen Tests | SaskHealthAuthority](#)
- [Virus Respiratory \(Combined Throat and Both Nostrils\) Specimen Collection Instructions | Public Health Ontario](#)

June 9, 2022

Public Health New Brunswick

4.11. Paxlovid Tool (for use in ARFs)

Pre-outbreak considerations: Is My Resident Eligible for Paxlovid (antiviral used to treat Covid 19)?

Paxlovid™ is an anti-viral medication intended for those who are at risk to experience more severe illness from COVID-19. There are many criteria that must be met before it can be prescribed. This is not an official eligibility assessment for Paxlovid, but merely a tool to help your facility to be ready for the process.

Complete the following pre-outbreak considerations for each resident in your facility. A resident who has all “Yes” responses is better prepared for a PAXLOVID assessment, should they test positive for COVID-19.

Pre-outbreak considerations:	Yes	No
1) Can the resident swallow full size pills without crushing? A resident must be able to swallow pills whole to qualify for PAXLOVID.		
2) Does the family/resident consent to resident receiving PAXLOVID, if they qualify? Informed consent of family/resident is required. For more information, review Paxlovid™ Patient Information		
3) Is the resident at risk to experience more severe illness from COVID-19? People at risk to develop more serious forms of Covid 19 include: <ul style="list-style-type: none"> • ≥ 18 and at higher risk for severe outcomes (see list) 		
4) Is the resident’s medication list up to date? Some medications cannot be used in combination with PAXLOVID. Contact your attending pharmacist to get a complete list of medication ready for each resident.		
5) Does the resident have a personal account for eVisitNB and MyHealthNB? <ul style="list-style-type: none"> • PCR test results can be obtained faster using MyHealthNB. • If unable to contact a resident’s health care provider, a prescription for Paxlovid may be obtained by contacting participating pharmacy, 811 or eVisitNB. *eVisit is Free until March 2023 with valid New Brunswick Medicare card. For access to this FREE service, please ensure you’re assessed by eVisit and not another online service company (EX. Maple) 		

If your resident has obtained YES responses for questions 1-4, they may be eligible for Paxlovid. Facility operators can further their readiness by taking the steps outlined in questions 5.

HOW TO OBTAIN A PAXLOVID ASSESSMENT FOR RESIDENTS

1. A resident in your facility has [symptoms of COVID-19](#). Have the resident’s symptoms started within the last 5 days?

Paxlovid must be started within 5 days of symptom onset. It is intended for people experiencing mild-moderate illness. If symptoms are severe, they DO NOT qualify for PAXLOVID. For a complete list of COVID symptoms go to: [About COVID-19 \(gnb.ca\)](#)

2. Test to confirm COVID-19:

- To access Paxlovid, individuals must have a confirmed positive test, either PCR or Point of Care Test (POCT). If a POCT is used, it is recommended to be observed by a health care provider or done by staff trained in POCT. ***Note: Contact EMP if concerned about health of resident.**

If test result is positive for COVID-19, resident may be eligible for Paxlovid.

If resident is followed by EMP:	If resident is not followed by EMP but has a primary care provider.	If resident is NOT followed by EMP AND does NOT have a primary care provider (or cannot reach their primary care provider).
Contact EMP. EMP will follow-up with Primary Care Provider to determine if resident is eligible for PAXLOVID and will send completed eligibility form to your attending pharmacy.	Notify primary care provider Primary care provider will determine if resident is eligible for PAXLOVID and will send completed eligibility form to your attending pharmacy. If unable to connect with their physician/nurse practitioner, contact a participating pharmacy to be assessed. List of participating pharmacies HERE	Contact a participating pharmacy to be assessed. (List of participating pharmacies HERE) If unable to be assessed by participating pharmacy, contact 811 or visit www.eVisitnb.ca*. Get an online New Brunswick Physician/Nurse Practitioner to assess resident’s eligibility for PAXLOVID. For access to this free service, please ensure you’re assessed by eVisit and not another online service company (Ex. Maple)
*Note: Contact EMP if concerned about health of resident or need support accessing Paxlovid.		

ONCE A PRESCRIPTION IS OBTAINED:

3. Obtain PAXLOVID from your attending pharmacy for eligible residents. Carefully follow instructions provided with this medication. Additional resources: Refer to [LIVING WITH COVID: management of COVID-19 for NB Long Term Care Homes and Caring for someone with COVID-19 \(gnb.ca\)](#)

Resident has symptoms of COVID-19
 Refer to [LIVING WITH COVID: management of COVID-19 for NB Long Term Care Home](#) and [Caring for someone with COVID-19 \(nbc.ca\)](#)

To access Paxlovid, individuals must have a confirmed positive test, either PCR or Point of Care Test (POCT). If a POCT is used, it is recommended to be observed by a health care provider or done by staff trained in POCT. *Note: Contact the EMP Care Coordination Center at 1-844-982-7367 if concerned about health of resident.

- TIPs to be prepared:**
- ✓ Make sure medication lists are up to date.
 - ✓ Provide Paxlovid information to resident or next of kin
 - ✓ Create a [MyHealthNB](#) account for each resident

COVID-19 Test results?

Negative:
 Re-test if new or worsening symptoms appear and/or contact Primary Care Provider for further investigation.

Positive. Your resident may be eligible for Paxlovid.
 If resident is yes to all the following, proceed for Paxlovid assessment by Physician/Nurse Practitioner

- ✓ Age: \geq 18 and at higher risk for severe outcomes (see list)
- ✓ resident's symptoms started within the last 5 days
- ✓ resident can swallow full size pills without crushing
- ✓ family/resident consent to resident receiving PAXLOVID if they qualify [Paxlovid™ Patient Information](#)

Is your resident followed by EMP?

NO

YES

Contact local EMP Unit. They will contact Physician/Nurse Practitioner to assess resident for PAXLOVID eligibility.

Does your resident have a Physician or Nurse Practitioner?

NO

YES

Contact a participating pharmacy to be assessed. (List of participating pharmacies [HERE](#))
 If unable to be assessed by participating pharmacy, contact 811 or visit [www.evisitnb.ca](#)*. Get an online New Brunswick Physician/Nurse Practitioner to assess resident's eligibility for PAXLOVID.

Contact EMP Care Coordination Center if there are concerns regarding health and/or need support accessing Paxlovid

Notify their physician/nurse practitioner to assess resident's eligibility for PAXLOVID. If unable to connect with their physician/nurse practitioner, contact a participating pharmacy to be assessed.

*NOTE: Each resident will need an account to access eVisit. This service will be free until March 2023 with a valid New Brunswicker Medicare card. For access to this free service, please ensure you're assessed by eVisit and not another online service company (Ex. Maple).